


U. S. DEPARTMENT OF AGRICULTURE

FARMERS' BULLETIN No. 1458

July 1930

STRAWBERRY DISEASES



STRAWBERRY DISEASES

By NEIL E. STEVENS, *Senior Pathologist, Office of Horticultural Crops and Diseases, Bureau of Plant Industry*

CONTENTS

	Page		Page
Introduction.....	1	Diseases of which causes are not definitely known.....	11
Leaf diseases.....	2	A leaf disease.....	11
Leaf spot.....	2	Root rots.....	12
Scorch.....	2	Fruit rots.....	12
Control of leaf spot and scorch.....	3	Gray-mold rot.....	12
Leaf blight.....	4	Tan rot.....	13
Mildew.....	5	Leather rot.....	13
Diseases caused by nematodes.....	5	Hard rot.....	14
Nematode gall.....	5	Fruit rots in transit and on the market.....	14
Root knot.....	6	Leak.....	14
Dwarf.....	6	Control of fruit rots.....	15
Virus diseases.....	8		
Strawberry yellows.....	8		
Witches'-broom.....	10		
Control of virus diseases.....	10		

INTRODUCTION

STRAWBERRIES are more widely grown in the United States than any other small fruit. Not only is the strawberry particularly adapted to the home garden and found to some extent in such gardens in every State, but centers of commercial production are located in many widely different regions throughout the country. From these large centers of production fresh fruit is distributed over a period of more than six months each year. The growth of this fruit in so many different regions under widely different conditions has resulted in the development of special cultural methods particularly adapted to local requirements, and the locations of the growing centers with reference to the markets served have given rise to somewhat different handling and shipping practices.

The number of varieties of strawberries grown in this country is large. Although about 50 varieties may be considered of commercial importance, experimental testing plots often contain several hundred. New varieties are grown from seed with ease, and plants of several new varieties are offered to the trade each season.

In view of the extent of strawberry culture in this country, the different conditions under which the crop is grown, and the number of varieties cultivated, it is obvious that strawberry diseases will vary in importance in different sections. Consequently, few general statements as to the severity of a given disease or the resistance to disease of strawberry varieties can be made with the expectation that they will be everywhere equally applicable. Methods of control must also be adapted to local conditions. What is undertaken in this bulletin is to describe the more important strawberry diseases and to give an account of such control methods as are of proved value.

LEAF DISEASES

The diseases here grouped under the general term "leaf diseases" will sometimes be found to affect other parts of the plant. Leaf spot and scorch often occur on fruitstalks and mildew on the young fruit. They are most common and conspicuous, however, on the leaves.

LEAF SPOT

One of the most widely distributed and best-known diseases of strawberries, leaf spot,¹ as its name indicates, is usually confined to the leaves. When the spots caused by the disease first appear they are of a reddish or purplish tint, but as they increase in size the center of each becomes paler and finally gray or almost white. A typical fully developed spot is about an eighth of an inch in




FIGURE 1.—Strawberry leaf spot. (Photographed by Wolf)

diameter and has a dead white center surrounded by a distinct reddish or purplish border, which finally merges into the color of the healthy leaf. (Fig. 1.) The spots are scattered irregularly over the surface of the leaf and often become so numerous as to cause the death of a large portion of the leaf and in extreme cases even of the plant. Usually, however, the injury due to this disease is chiefly that of weakening the plant and reducing the crop through the loss of needed leaf surface.

SCORCH

Scorch² is almost as common on strawberries in this country as leaf spot, for which disease it may be mistaken in the early stages,

¹ Caused by *Mycosphaerella fragariae* (Tul.) Lindau.

² Caused by *Mollisia earliana* (E. and E.) Sacc. (*Diplocarpon earliana* (E. and E.) Wolf).

when the spots or blotches, usually purplish in color, are irregular in outline and vary in diameter from one-twentieth to one-fifth of an inch. (Fig. 2.)

The diseased areas gradually enlarge, and many of them grow together. In severe cases the whole leaf may become purplish or reddish in color. Soon, however, the leaf margins become dry (fig. 3) and have the burned or scorched appearance which has given rise to the name of the disease.

In case of severe infection entire plants may be killed or so weakened as to be worthless. One type of injury caused by this disease that results in large losses is the infection which often occurs on the fruitstalks, sometimes girdling them and thus causing the death of the flowers and young fruit.




FIGURE 2.—The "spot" stage of strawberry scorch

CONTROL OF LEAF SPOT AND SCORCH

That leaf spot can be controlled by spraying with Bordeaux mixture has been known for many years. Experiments by the Connecticut and North Carolina Agricultural Experiment Stations have demonstrated that scorch may also be controlled by this method.

It has been found most satisfactory for the control of leaf spot and scorch to spray with 4-4-50 Bordeaux mixture as soon as growth is well started in the spring—that is, when the first leaves are about half grown—and to repeat the treatment as often as may be necessary to keep the foliage well covered until the first berries are about one-third grown. It may be necessary to spray as often as every 10 days. Usually one or two additional applications in the late summer or fall will result in much better control.

Whether spraying should be undertaken in any particular case must be decided by the individual grower, taking into consideration

its cost and the probable extent of the loss from these diseases. Often it proves more economical merely to reduce the diseases so far as is practicable by frequent renewal of strawberry patches and by reducing the chances for infection. Mowing off and burning old leaves and removing diseased leaves before plants are set tend to reduce infection from strawberry scorch and leaf spot.

It is wise also to avoid setting any varieties that appear particularly susceptible to these diseases in a given locality. Lists of strawberry varieties that have proved very susceptible or resistant to certain diseases have been published at various times. These come from different workers and are frequently contradictory, the fact being, of course, that a variety may be obviously susceptible in one place or in one season and almost free from disease in another place or during another year. Reliable information as to the relative sus-




FIGURE 3.—A more advanced stage of strawberry scorch. The leaves are beginning to dry about the margins. (Photographed by Wolf)

ceptibility of different varieties to various diseases is very difficult to obtain, owing to the fact that in large strawberry-growing sections one or two varieties are likely to be grown to the exclusion of others, and experimental plots usually contain relatively few plants of any given variety and are often maintained for only a few years. In order to secure reliable information on this point it would be necessary to carry on extensive comparative tests of the leading varieties in various sections of the country for a period of at least five or six years. The wisest policy for the present will be for each grower to select varieties on the basis of local experience in the region.

LEAF BLIGHT

Another disease which causes injury to strawberry leaves and leaf-stalks and has been found to be serious in Illinois and Michigan is

angular spot, sometimes called blight.³ The dead areas caused by this disease even in the early stages are much larger than those made by leaf spot. Usually there is only one injured region to a leaflet, although there may be three or four. In an advanced stage this disease is distinguished by the more or less triangular shape of the dead area, which often extends from one of the larger veins to the side of the leaflet and includes all the leaf tissue between the edge of the leaflet and the vein.

MILDEW

Mildew⁴ has long been recognized as a serious disease in England and is locally destructive in the United States. The most conspicuous symptom of the disease is the curling of the leaves. The edges of the leaflets gradually turn upward until much of the under side is exposed. If the under side of a diseased leaf is closely examined it will be found to be covered with a whitish down composed of the causal fungus.

For the commercial control of strawberry mildew in this country the ordinary sanitary measures recommended for leaf spot and scorch have usually proved sufficient. However, the New York (Geneva) Agricultural Experiment Station has demonstrated in the Hudson Valley that mildew may be practically eliminated even during seasons of severe infection by four applications of 85-15 lime-copper sulphate dust. The first application was made as the first buds were expanding in the cluster and the succeeding ones at intervals of about 12 days.

DISEASES CAUSED BY NEMATODES

Three distinct diseases of strawberries, namely, root knot, gall, and dwarf, are caused by nematodes. These are wormlike animals, and the species that affect strawberries are very small—from one-fiftieth to one twenty-fifth of an inch in length. The nematode gall of strawberries has been found only in the Northwest. Root knot, on the other hand, is found to some extent in all except the more northern States, but is usually serious on strawberries only in the Southern States. Dwarf is common in the Southeastern States.

NEMATODE GALL

The disease of strawberries known as gall⁵ was first found in the United States in Oregon and is known only in the northwestern part of the country. The disease appears to affect only those portions of the plant which are above the ground. The plants as a whole are usually somewhat dwarfed and their color is of a lighter green than that of normal plants. As shown in Figure 4, affected leaves become wrinkled, swollen, and often much reduced in size. On the leaf and fruit stems are often formed conspicuous galls which contain the nematodes in great numbers. The nematode which causes gall on strawberries is known to occur in the United States on hyacinth, red clover, alfalfa, false dandelion, and the wild strawberry (*Fragaria chiloensis*) of the Pacific coast. It has been

³ Caused by *Dendrophoma obscurans* (E. and E.) H. W. And.

⁴ Caused by *Sphaerotheca humuli* (DC.) Burr.

⁵ Caused by the nematode *Tylenchus dipsaci* (Kühn) Bastian.

proved by inoculation experiments that nematodes from strawberries will affect red-clover seedlings. This suggests the possibility that the pest passes from one host to another in nature and that it may be dangerous to plant strawberries in fields in which affected crops of red clover or alfalfa have been grown. For the present, however, the most necessary control measure is the obvious one of not planting strawberries that are infected or that come from infested fields.

ROOT KNOT

As indicated by its name, the disease known as root knot^a is characterized by swellings or enlargements of the roots. Usually root knot produces no malformations on those parts of the plant which are above ground, and as a result it is frequently overlooked. If the roots of an infected plant are examined, however, the disease is usually readily detected. In severe cases great numbers of the finer roots and many of the large roots are found to be much enlarged at various points. These swellings, which are caused by the nematodes within the roots, vary greatly in size and shape.

The extent of injury to strawberries due to root knot is very difficult to estimate. Relatively few strawberry fields in the older growing regions of the South are free from root knot, and the reduced yield due to the disease has come to be taken as a matter of course. Often, however, the disease becomes serious enough to cause the death of part of the plants and can no longer escape notice.

When available, new land or land known to be free from the root-knot nematode should be used for strawberries. Infested fields, however, may be largely freed of infestation by a three-year rotation with crops which are immune to this nematode.

The following plants are known to be immune or only slightly susceptible to root knot:

Barley.	Grasses (nearly all).	Sorghum.
Beggarweed.	Kafir.	Soybean (Laredo variety).
Broomcorn millet.	Milo.	Timothy.
Chufa.	Natal grass.	Velvetbean.
Corn.	Peanut.	Wheat.
Cowpea (Brabham, Iron,	Pearl millet.	Winter oats.
Monetta, and Victor varieties).	Redtop.	
	Rye.	

DWARF

The most conspicuous disease of strawberry plants in the southeastern part of the United States during the summer months is the one here referred to as dwarf, by which name it is generally known to growers in North Carolina, Louisiana, and elsewhere. In Florida, where the disease has long been known, it has received various names, including crimps, French bud, white bud, and brier bud.

The most conspicuous symptom of the disease and the one from which all its common names are derived is the malformation it induces in the young leaves as they develop. The leaves of diseased plants become greatly reduced in size and much deformed. (Fig. 5.) The

^a Caused by the nematode *Caenorhynchus radiculicola* (Greef) Cobb. A full description of the root-knot nematode, its characteristics on various hosts, and the known methods of control are given in Farmers' Bulletin 1345.

edges of the leaflets are crinkled. In extreme cases the entire plant is dwarfed and the internodes of the runners are shortened. The disease is apparently abundant throughout the southeastern part of the United States from Virginia to Louisiana. It is being actively



FIGURE 4.—Nematode gall of strawberry. (Photographed by Byars)

studied by investigators in several States, who agree upon the following points:

The disease is caused by a nematode.⁷

The symptoms are most conspicuous during the summer months, and diseased plants are often hard to distinguish in winter and spring.

⁷*Aphelenchus fragariae* Ritzema-Bos or a closely related form.

The disease is often introduced into new fields by the planting of infected plants, and spreads to healthy plants.

The disease is often spread by water and is most common in the low portions of the fields.

Runner plants from dwarf mother plants are usually dwarfed.

The disease is very common on strawberries of the Klondike and Missionary varieties and has recently been found on the Blakemore, proving that this valuable new variety is not immune to it.




FIGURE 5.—Typical "dwarf" strawberry plant of the Klondike variety. (Photographed by Mook at Chadbourn, N. C., August, 1928)

When strawberries are hoed any plants showing symptoms of dwarf should be removed.

VIRUS DISEASES

STRAWBERRY YELLOWS

Yellows is common on strawberries of the Marshall (Banner) variety in central California. The most conspicuous symptoms of the disease are (1) characteristic curling and cupping of the leaves, (2)

yellowing of the leaves around the margin between the larger veins, and (3) dwarfing of the leaves and a general stunting of the growth of the entire plant. (Figs. 6 and 7.)



FIGURE 6.—Young Marshall (Banner) strawberry plant showing typical yellows. (Photographed by Plakidas)

This disease is transmitted through the runners; that is, every runner plant from a diseased mother plant will be diseased. It is not transmitted through seed. The root system, however, appears healthy during the early stages of the disease. Under favorable conditions for growth the plants sometimes appear to recover their

vigor, but later show the symptoms of disease again. Complete recovery of the plant has not been noted.

Carefully controlled experiments, however, have proved that it is transmitted by a species of strawberry aphid (*Myzus fragaefolii* Cockerell), which fact, together with other characteristics, would indicate that it belongs to the class of virus diseases.

WITCHES'-BROOM

A disease of strawberries which seems to be of the virus type was found in western Oregon in 1925 and is known only from this region. This disease, which is transmitted by the strawberry leaf aphid (*Myzus fragaefolii*), was named witches'-broom because one of the most characteristic symptoms of the disease is that the leaf petioles



FIGURE 7.—Sister seedling plants of Marshall (Banner) variety of strawberry. Left, diseased with yellows; right, healthy. (Photographed by Plakidas)

are more slender than normal ones and the leaves tend to bunch together. The runners of diseased plants become shortened so that runner plants stand close to the mother plants. The leaflets are lighter in color than in normal plants, and the margins arch downward. The "broom" character is exhibited more by some varieties like Ettersburg No. 121 than by others. (Fig. 8.)

CONTROL OF VIRUS DISEASES

So far as our present knowledge goes, control of virus diseases in strawberries must depend on the selection of healthy stock and on the elimination of diseased plants as they appear. Since strawberries are usually grown with a short rotation period and much handwork is usually involved in their culture, the destruction of diseased plants at frequent intervals should be readily accomplished.

DISEASES OF WHICH CAUSES ARE NOT DEFINITELY KNOWN

A LEAF DISEASE

A disease which resembles a mosaic in appearance and which has thus far been found only in the northeastern part of the United States and in Canada is undoubtedly different both from the yellows of the Pacific coast and from the dwarf of the Southern States. Yellowing is the most striking symptom. The yellow color is so prominent that the diseased plants in a field may easily be seen from a distance. In extreme cases as much as 90 per cent of the area of a leaf may be devoid of chlorophyll. The yellow color is




FIGURE 8.—Strawberry plant of the Ettersburg No. 121 variety affected with witches'-broom in which the runner plants have been pulled away from the mother plant to show the shortened internodes of the runners. (Photographed by Zeller)

usually of a light hue, nearly albino. The affected leaves usually show a mosaiclike pattern of mottling or marbling, a fact which has led many of those who have seen the disease in the field to designate it as mosaic. The yellow leaves appear thinner than the healthy green ones. Curling, crinkling, cupping, or twisting of the leaflets is rare. The symptoms are most prominent during the cool weather of spring. The yellowing practically disappears in late summer. The disease seems to be of a systemic nature; that is, all runner plants of a diseased mother plant are affected. The appearance of the disease and its behavior in the field suggest that it is of the virus type, but as yet transmission experiments have uniformly given negative results.

ROOT ROTS

Species of *Rhizoctonia* have been reported as associated with "black root" in Michigan, Tennessee, Washington, and other States. In Canada species of *Fusarium* and certain soil bacteria have been found associated with this type of disease, and a species of *Diplodina* was found apparently causing a disease of strawberry roots in England.

In Lanarkshire, Scotland, root diseases of strawberries have become very important in recent years. A species of *Pythium* has been isolated from diseased roots and proved by inoculation to be capable of infecting healthy plants. The work in that locality indicates, however, that while this and perhaps other fungi are no doubt an important factor in the injury, the condition of the soil and the general health of the plant are also of great importance.

In the writer's work on root diseases of strawberries, species of *Rhizoctonia*, *Chaetomium*, *Thielavia*, *Melanospora*, and *Hypocopa* appear most commonly associated with strawberry roots.

During recent years strawberry growers have sustained heavy losses in various parts of the country from what appear to be root diseases. These diseases, which have usually been designated as "black root" or "red root," have been studied by investigators in various States, and several fungi have been isolated from the diseased roots. Thus far, however, all attempts at the production of disease by inoculation with these fungi have failed. This failure by no means proves that these or other fungi are not responsible for the trouble, and certain fungi found on strawberry roots are now being actively studied. Meanwhile no information is available on which to base recommendations for control. For the present, growers must depend for control on selection of stock, careful culture, and frequent rotation, the last probably being the most important.

FRUIT ROTS

Fruit rots of strawberries, both on the plant and after the berries are picked, are caused by a variety of diseases, some of which are now known only in certain localities. In discussing each disease the region in which it is known to be serious will be indicated. This does not mean, of course, that the disease does not occur in other places; indeed, it is highly probable that many of these rots will be found in regions widely separated from those in which they are now known.

GRAY-MOLD ROT

The best known and most widely distributed fruit rot of strawberries is that caused by gray mold (*Botrytis* sp.). This disease affects both green and ripe berries. The rot usually starts in the part of the berry that is next to the ground or that touches another berry or a leaf. It is first noticed as a light-brown rather soft spot. The rot spreads in all directions through the berry and is not localized in any particular portion. After the berry is wholly rotten it soon begins to dry out and becomes firm and tough and of a uniform color throughout. At this stage the berry becomes almost entirely covered by a firm gray powder or dusty appearing fungous growth,

the well-known gray mold. This disease is most common in the cooler strawberry-growing regions. It is often very destructive in Maryland, Delaware, New York, and New England, where it is not unusual for 10 per cent or more of the crop to be lost. The disease is especially favored by moisture and may become destructive in any of the large strawberry-growing sections during continued wet weather.

The growth of gray mold can not be entirely prevented by refrigeration. Under the conditions found in many refrigerator cars, strawberries affected with this disease often become covered with an abundant growth of soft, fluffy, gray mold, which is very conspicuous and seriously hurts the sale of the entire load. Consequently, it is important that berries affected with gray mold be carefully culled out during picking and packing.

The injury caused by gray mold is not confined to the fruit. The blossoms are frequently blighted by this disease, and under favorable conditions leaf and flower stalks may be infected and the whole plant thus seriously injured.

TAN ROT

The so-called tan rot⁸ is important, especially during rainy seasons, in Louisiana, Florida, Arkansas, Virginia, Maryland, and North Carolina. It may attack either green or ripe berries, forming slightly sunken tan-colored areas, which usually develop on the side of the berry. The rot extends into the pulp so that the rotten portion forms a cone with its base at the surface of the berry. The outer layer of this diseased area becomes soft and as the core of the rotten portion is held together by the rot fungus it is easily removed intact with the point of a knife or pencil. This characteristic serves to identify this trouble readily.

LEATHER ROT

Leather rot,⁹ which has been found in Alabama, Louisiana, Mississippi, Arkansas, Missouri, Tennessee, Kentucky, Illinois, Maryland, and Virginia, is readily distinguished from other fruit rots by the bitter taste of berries even slightly rotted. This disease may affect either ripe berries or green ones in any stage of development. When ripe berries are affected, all parts become dark brown over the affected areas or brown at the edges, shading off into the natural green. Partly colored berries show the same light-brown color at the center, but the spot shades off into purple and finally to the red of the berries. If a berry affected with this disease is cut across, a marked browning of the vascular strands is readily noted. This is usually accompanied by a less marked browning of the other tissues. At no time is there any clear line separating the diseased and healthy flesh, nor can the diseased portion be easily removed, as in the case of tan rot. The disease is most noticeable after rainy periods, especially if the weather is warm. Leather rot has proved very serious during recent years in White County, Ark., and in Tennessee. During some years more than one-fourth of the crop in these localities has been lost.

⁸ Caused by *Peizizella lythri* (Desm.) Shear and Dodge.

⁹ Caused by *Phytophthora cactorum* Leb. and Cohn.

HARD ROT

Hard rot¹⁰ is known to be serious in the strawberry fields of central Florida, and it has been found in North Carolina, Arkansas, and Tennessee. As the rot is caused by a soil fungus, it almost always occurs on the side of the berry that comes in contact with the soil, and a small quantity of soil will usually be found clinging to the decayed spot.

The affected berries are generally one-sided and show a hard brown rot separated from the sound portion of the berry by a distinct line. The portion of the berry not actually penetrated by the rot fungus remains unchanged both in appearance and in taste.

FRUIT ROTS IN TRANSIT AND ON THE MARKET

The various rots just described are all found to a greater or less extent on berries in the field. A very large proportion of the rotten berries in most cull piles are infected with one or more of the fungi just mentioned. Their relative abundance varies in different parts of the country and with weather conditions. In general, as every strawberry grower knows, these field rots are more abundant during or immediately after a period of wet weather. It is during such rainy periods that the greatest number of rotten berries must be thrown out as culls when the berries are sorted in the packing shed.

Serious as is the loss from field rot to the grower, it is not as costly as the rots that occur in transit and on the market. Field rot as such means a reduction in the size of the crop and some added cost in the packing shed. Rot that develops in transit, however, means that the labor of growing that much of the crop has been wasted, as well as the cost of picking, packing, and hauling to the railroad. To this waste must be added the expense of loading in the cars and the cost of freight or express and of refrigeration. Often more important than any of these is the disastrous effect on the market of the presence of moldy or leaking berries in the crates that are offered for sale.

All of the fruit rots described above are known to develop to some extent in transit. Gray mold, because of its wide distribution and its ability to grow somewhat at the temperatures usually obtained in refrigerator cars, is the most important in transit of any of the brown rots.

More important than any of the brown rots, indeed more important in transit and on the market than all of them put together, is the rot almost universally known as "leak."

LEAK

As its common name indicates, leak¹¹ is characterized by the collapse of the berry and the loss of much juice. Even one or two leaking berries will stain a box and make it unattractive, and it is not uncommon for the juice to drip from boxes and crates of badly infected fruits. Spores of the black mold are almost always abundant in the air of packing sheds, stores, and houses. Once inside the

¹⁰ Caused by *Rhizoctonia* sp.

¹¹ Caused by the common black mold, *Rhizopus nigricans* Ehrh.

fruit, black mold will destroy a strawberry much more quickly than any other fungus now known. Fortunately, however, this fungus usually enters the berry only through wounds and grows very slowly at temperatures below 50° F. Control measures, then, must be directed toward careful handling and the maintenance of low temperatures, as will be indicated in the following sections.

The importance of leak as a cause of loss in strawberries is strikingly shown by the inspection certificates of the food products inspection service of the Bureau of Agricultural Economics, United States Department of Agriculture. A careful compilation of the reports relating to strawberries examined during the 7-year period ended December 31, 1925, which included inspections of 5,370 carloads of strawberries, showed that the loss caused by leak (*Rhizopus* rot) was practically one-fourth greater than that caused by all other rots combined, the exact figures for the amount of loss being 4.9 per cent for leak and 3.9 per cent for all others. In certain years and in certain localities other rots often exceed *Rhizopus* rot, but for the whole country and over this period *Rhizopus* rot was more important than all of the other rots put together. This same report also shows in an interesting manner the difference between the amount




FIGURE 9.—Percentage of *Rhizopus* rot and of all other rots in carload shipments of strawberries from Arkansas, Tennessee, and Louisiana for two wet years (1922 and 1923) and two dry years (1924 and 1925)

of rot in wet and dry seasons. Figure 9 gives a comparison of the rot in Arkansas, Tennessee, and Louisiana during two wet years (1922 and 1923) and two dry years (1924 and 1925).

CONTROL OF FRUIT ROTS

The methods that have proved so successful in controlling fruit rots in grapes and most tree fruits, namely, spraying or dusting, have not proved practicable with strawberries. While experiments carried out in Arkansas did indicate that the amount of field and storage rot could be reduced in certain years by spraying or dusting, the rot proved not to be of sufficiently regular occurrence to justify the expense of treatment. There is no indication that any practicable application of spray or dust will reduce the loss from leak, which, as already noted, is the most important of all strawberry rots.

MULCHING

Mulching, if carefully done, reduces the loss from rots in two ways: (1) It reduces the infection from certain fungi which apparently gain entrance from the soil, especially leather rot and hard rot, and

(2) by keeping the berries free from soil it reduces the chance of bruising and thus lessens the opportunity for infection by black mold.

CAREFUL HANDLING

As has been proved to be the case with so many fruits, careful handling is one of the best means of reducing the loss from rot in strawberries. Careful handling is important at every stage—in picking, in sorting, in packing, and in loading the crates on trucks or into cars. This means the careful supervision of all employees and careful training of inexperienced pickers.

KEEPING THE BERRIES COOL

All rot fungi are favored by high temperatures. Although gray mold can not be entirely checked by any degree of refrigeration which can be safely secured at present, even this rot grows much more slowly at low temperatures. It is therefore of first importance that strawberries be kept as cool as possible from the time they are picked until the time they are eaten. A useful aid in accomplishing this and one that has proved most advantageous in practice is to pick as large a part of the crop as possible early in the day while the fruit is still cool. Not only does this result in the berries reaching the car cooler and thus requiring less ice to remove the "field heat," but the berries are actually less easily bruised if picked while cool.

Careful tests of the pressure required to puncture strawberries and other small fruits showed that the pressure required to puncture cool fruit was greater than that required to puncture warm fruit. This was equally true whether the fruit was wet or dry.

BEST PRACTICE WHEN STRAWBERRIES ARE WASHED BEFORE PACKING

In some sections, notably central Florida, cultural and trade practices make it necessary to wash the berries before they are packed. When this is done the berries should not be dried after washing but merely allowed to drain for a short time and packed while still moist, and kept in the shade until they are delivered to the loading platform. It is important that the water in which the berries are washed should be changed frequently. Dirty water not only contains sand, which makes it much easier to injure the berries in handling, but it contains different spores of mold fungi, particularly *Rhizopus*, which greatly increase the loss from decay.

REFRIGERATION IN TRANSIT

Careful handling to prevent bruising, and quick cooling after harvesting, are both important, but, of course, do not assure fruit reaching destination without loss. It is of first importance that the fruit be properly refrigerated in transit.

It is desirable to keep the fruit temperature in transit below 45° F.; 40° is still better. These temperatures, however, are rather difficult to maintain unless salt is used in the bunkers with the ice. For this purpose, at the initial icing the quantity of salt added should be from 1½ to 2 per cent of the ice by weight. At the first

reicing salt may be again added in the same proportion, that is, between $1\frac{1}{2}$ and 2 per cent of the additional ice.

A great aid in maintaining low fruit temperatures in transit is to precool the fruit after it has been loaded in the car and before the shipment moves to market. This may be done in various ways, one of the simplest of which is by the use of a precooling device developed by the United States Department of Agriculture.¹²

¹² Copies of a mimeographed circular describing this device and giving instructions for its use are available for distribution on request.

ORGANIZATION OF THE UNITED STATES DEPARTMENT OF AGRICULTURE

June 25, 1930

<i>Secretary of Agriculture</i> -----	ARTHUR M. HYDE.
<i>Assistant Secretary</i> -----	R. W. DUNLAP.
<i>Director of Scientific Work</i> -----	A. F. WOODS.
<i>Director of Regulatory Work</i> -----	WALTER G. CAMPBELL.
<i>Director of Extension Work</i> -----	C. W. WARBURTON.
<i>Director of Personnel and Business Administration.</i>	W. W. STOCKBERGER.
<i>Director of Information</i> -----	M. S. EISENHOWER.
<i>Solicitor</i> -----	E. L. MARSHALL.
<i>Weather Bureau</i> -----	CHARLES F. MARVIN, <i>Chief.</i>
<i>Bureau of Animal Industry</i> -----	JOHN R. MOHLER, <i>Chief.</i>
<i>Bureau of Dairy Industry</i> -----	O. E. REED, <i>Chief.</i>
<i>Bureau of Plant Industry</i> -----	WILLIAM A. TAYLOR, <i>Chief.</i>
<i>Forest Service</i> -----	R. Y. STUART, <i>Chief.</i>
<i>Bureau of Chemistry and Soils</i> -----	H. G. KNIGHT, <i>Chief.</i>
<i>Bureau of Entomology</i> -----	C. L. MARLATT, <i>Chief.</i>
<i>Bureau of Biological Survey</i> -----	PAUL G. REDINGTON, <i>Chief.</i>
<i>Bureau of Public Roads</i> -----	THOMAS H. MACDONALD, <i>Chief.</i>
<i>Bureau of Agricultural Economics</i> -----	NILS A. OLSEN, <i>Chief.</i>
<i>Bureau of Home Economics</i> -----	LOUISE STANLEY, <i>Chief.</i>
<i>Plant Quarantine and Control Administration</i> ..	LEE A. STRONG, <i>Chief.</i>
<i>Grain Futures Administration</i> -----	J. W. T. DUVEL, <i>Chief.</i>
<i>Food, Drug, and Insecticide Administration</i> ..	WALTER G. CAMPBELL, <i>Director of</i> <i>Regulatory Work, in Charge.</i>
<i>Office of Experiment Stations</i> -----	-----, <i>Chief.</i>
<i>Office of Cooperative Extension Work</i> -----	C. B. SMITH, <i>Chief.</i>
<i>Library</i> -----	CLARIBEL R. BARNETT, <i>Librarian.</i>